



Study of *Argemone mexicana* and *Candida albicans*

Lalit Rawat and Ragini Shukla

P. K. College, Contai, Purba Medinipur, Rajasthan, India

Abstract: Plants measure wealthy supply of medication for antifungal and antiviral properties. These properties are employed by plants to safeguard themselves from foreign particles or pathogens. With the advancement of techniques currently we tend to carry out extracting the chemicals and victimisation to cure plant and animal infection. The *in vitro* antifungal activity of binary compound extract from *Argemone mexicana* plants utilized in ancient drugs for the treatment of varied diseases.

Keywords: Antifungal, *Argemone*

I. INTRODUCTION

A variety of microorganisms as well as fungi, bacteria, viruses, harbour the soil. Out of those microorganisms, few area units useful for the soil, some don't damage the soil in any respect neither have the benefit of it, however a number of them have the potential to cause major injury to plant growth moreover because the quality of soil. Soil borne illness will be a serious limitation for plants. The infective agent might stay dormant within the soil and may become morbidic once more as shortly because the host is out there. Preventive measures taken for the soil to avoid contamination and healthy environmental conditions will be effective for healthy plant growth. Fungi are indirectly to be blamed for allergic or cytotoxic disorders due to the assembly of mycotoxins or allergens. Generally, for the management of phytopathogenic fungi use of artificial fungicides is finished. These chemical fungicides have tremendous effects on the plant as well as on the animals feeding on them Hence, there has been an increasing demand to create use of natural merchandise which will act as antifungal agents resulting in less injury to the surroundings and living organisms. Biologically active compounds found in plants are a lot safer than the artificial fungicides. Hence, extracts and oils of medicative plants has been used since it contains a great deal of secondary metabolites as compared to the other plants.

II. MATERIALS AND METHOD

2.1 Materials

A. Fungi Used

1. *Aspergillus niger*


Isolation: *Aspergillus* culture was obtained by doing agar plate method and the culture was observed under microscope and sub-culturing was done to obtain pure cultures of *Aspergillus niger*. Seven days old culture of fungi was used.

Macroscopic Characters: The colony is black in colour and the reverse surface is white to light yellow in colour.

Microscopic Characters: Hypha is septate and hyaline. Conidiophore is present and arises from the basal foot. Conidiophores are smooth-walled, hyaline or turning dark towards the vesicle.



Conidial heads are biserial with the phialides borne on brown, often septate metulae. Conidia 3.5-5.0 μm in diameter, dark brown to black and rough-walled.



2. *Candida albicans*

Isolation: *Candida* culture was obtained by doing agar plate method which gave white colonies. The colony was then sub-cultured and *Candida albicans* pure cultures were obtained. Seven days old culture of fungus was used.

Macroscopic Characters: Colonies are white to cream in colour.

Microscopic Characters: Abundant branched pseudohyphae and true hyphae with blastoconidia are present. The blastoconidia are formed in grape-like clusters along the length of the hyphae. Terminal chlamydoconidia may be formed with extended incubation.



B. Plants Used

Argemone mexicana:

Species of poppy found in Mexico and now widely naturalized in many parts of the world. It has bright yellow latex. It is poisonous to grazing animals, and it is rarely eaten, but it has been used medicinally by many peoples. *Argemone mexicana* seeds contain 22–36% of a pale yellow non-edible oil, called *argemone oil* or *katkar oil*, which contains the toxic alkaloids sanguinarine and dihydrosanguinarine. Four quaternary isoquinoline alkaloids, dehydrocorydalmine, jatrorrhizine, columbamine, and oxyberberine, have been isolated from the whole plant of *Argemone mexicana*. Many other alkaloids such as argemexicaines A and B, coptisine, cryptopine, allocryptopine and chelerythrine have also been found in this plant. The seed pods secrete a pale yellow latex when cut open. This argemone resin contains berberine and protopine.





C. Culture Used

1. Liquid Culture Medium: Potato Dextrose Broth

Potato: 200g, Dextrose: 20g, Distilled water: 1000ml

Peeled and chopped potatoes were boiled in distilled water, till the water became starchy. Solution was filtered through muslin cloth and the volume was raised to 1000 ml by adding distilled water. Filtrate was then transferred to a conical flask and dextrose was added. The flask was plugged and autoclaved at the pressure of 15 lbs./sq. inch at 121°C for 20 minutes. The pH of the media was checked (5.6 ± 0.2). Streptomycin was added before pouring the plates.

2. Solid Culture Medium: Potato Dextrose Agar

Potato: 200g, Dextrose: 20g, Distilled water: 1000ml, Agar: 14gm

Peeled and chopped potatoes were boiled in distilled water, till the water became starchy. Solution was filtered through muslin cloth and the volume was raised to 1000 ml by adding distilled water. Filtrate was then transferred to a conical flask and dextrose and agar were added. The flask was plugged and autoclaved at the pressure of 15 lbs./sq. inch at 121°C for 20 minutes. The pH of the media was checked (5.6 ± 0.2). Streptomycin was added before pouring the plates.

D. Other Requirements

Petri plates, Conical flasks, Micropipettes, cork borer, forceps, nichrome loop, Laminar Air Flow, etc. The entire experiment must be carried out in Aseptic conditions with sterilized glasswares.

2.2 Method

A. Aqueous extract preparation from *Argemone mexicana*:

Leaves were crushed and soaked in Distilled water and kept on shaker overnight (150 rpm for 24 hours). Next day the mixture was filtered using muslin cloth. The extract was stored at 4°C.

Extract of Different Concentration:

Concentration (%)	Dried powder (gm)	Distilled water(ml)
10%	10	100
20%	20	100
30%	30	100
40%	40	100
50%	50	100

**B. Testing of Plant aqueous Extract of *Argemone mexicana*:**

1. In Liquid Culture Medium: 49 ml of Potato dextrose Broth was added to each conical flask and 1 ml of plant extract was added to it. The total volume should be 50 ml in each conical flask. A control flask was maintained to compare the growth. The flasks were plugged and autoclaved for sterilization of Potato dextrose broth and extract.

Preparation of the concentrations of aqueous extract of *Argemone mexicana*:

Concentration g/ml (wt/vol)	Volume of extract(ml)	Volume of broth (ml)
Control	0	50
10	1	49
20	1	49
30	1	49
40	1	49
50	1	49

III. OBSERVATIONS AND RESULTS**1. In liquid culture medium:**

Control	Excellent	++++
10	Very Good	+++
20	Good	++
30	Fair	+
40	Poor Growth	+
50	No Growth	-

***Argemone mexicana* extract on *Aspergillus niger*:**

Concentration of Extract	Weight of Whatmann filter paper (g)	Weight of paper + Biomass (After drying)	Weight of Biomass (g)	Visual Fungal growth in PD Broth
Control	0.830	1.612	0.782	++++
10%	0.820	1.517	0.697	++++
20%	0.822	1.484	0.662	+++



30%	0.831	1.478	0.647	++
40%	0.821	1.440	0.619	+
50%	0.823	0.874	0.051	-



Aspergillus niger grown in PD Broth + *Argemone mexicana* aqueous extract

***Argemone mexicana* extract on *Candida*:**

Concentration of Extract	Weight of Whatmann filter paper (g)	Weight of paper + Biomass (After drying)	Weight of Biomass (g)	Visual Fungal growth in PD Broth
Control	0.831	1.809	0.978	+++++
10%	0.882	1.158	0.276	+++
20%	0.825	1.078	0.253	+++
30%	0.812	0.988	0.176	++
40%	0.822	0.895	0.073	++
50%	0.841	0.871	0.030	-



Candida albicans grown in PD Broth + *Argemone mexicana* aqueous extract

2. In Solid Culture Medium

Argemone mexicana extract on *Candida*

Concentration of Extract	Inhibition zone observed on the 8 th day (cm)
--------------------------	--



	I	II	Average
Control	0.2	0.6	0.4
10%	0.1	0.8	0.45
20%	0.1	0.2	0.15
30%	0.1	0.6	0.35
40%	0.1	0.2	0.15
50%	0.3	0.4	0.35

IV. RESULTS

Results showed that *Argemone mexicana* was effective as an antifungal extract since it showed Zone of Inhibition. Although, *Aspergillus niger* was resistant, still it did show reduction in biomass at higher concentrations of plant extracts. *Candida* on the other hand was very sensitive to plant extracts and hence grew slowly.

V. CONCLUSION

The above experiment has concluded that that Aqueous plant extract of *Argemone mexicana* is effective against the both pathogenic fungi i.e., *Aspergillus niger* and *Candida albicans*. *Aspergillus niger* showed more resistance at most concentrations. *Candida* being sensitive to the antifungal compounds present in the extracts shows considerable decrease in biomass and showed complete inhibition at the highest concentrations used (50%). The variations in the effects of plant extracts on inhibiting the pathogenic fungi was due to the variable concentrations of plant extract we used. Also, the of solvent used for plant extract preparation has an effect on inhibition. Studies indicate that the higher the concentration of plant extract the higher is effect of inhibition. Higher concentrations of the plant extract show tremendous reduction in *Aspergillus niger* growth and complete inhibition for *Candida*. *Aspergillus niger* does not show zones of inhibition even on higher concentrations whereas, *Candida* does.

REFERENCES

- [1]. A.K. Pandey, M.K. Rai & D. Acharya (2003) Chemical Composition and Antimycotic Activity of the Essential Oils of Corn Mint (*Mentha arvensis*) and Lemon Grass (*Cymbopogon flexuosus*) Against Human Pathogenic Fungi, *Pharmaceutical Biology*, 41:6,421-425
- [2]. Abhishek Mathur et al. *Adv. Appl. Sci. Res.*, 2011, 2 (2):260-264.
- [3]. Agrios, G. N. (1988). *Plant Pathology*, 3rd. ed. Academic Press, Inc.: New York. 803. As cited in: Sahi, I. Y., A. N. Khalid, 2007.
- [4]. Alexopoulos, C. J., C. W. Mims. (1979). *Introductory Mycology*. Third edition.
- [5]. Ambikapathy, V. et al. *Asian J. Plant Sci. Res.*, 2011, 1(3):131-34.
- [6]. Barnett, H. L. and B. B. Hunter. (1972). *Illustrated genera of imperfect fungi*.